

# Overactive bladder

## We can help you regain control of your bladder

Your bladder has to store enough urine (wee) (between 250 – 500 milliliters) to allow you to do daily activities and have enough sleep.

When your bladder is filling up, the bladder muscle is relaxed. When the bladder starts to become hold on until it is convenient to go. When we decide to go to the toilet, the bladder muscles begin to contract. Adults usually empty their bladder every three to four hours during the day.

At night it is normal to need the toilet once. If you are over 60 years old you will probably find you get up to urinate more than once and by the age of 90 you may need to get up four times a night.

## If you have an overactive bladder you may have some or all of the following symptoms:

- Urgency
- Reduce the number of times you get up during the night
- Helps to stop urge incontinence

Bladder retraining requires you to hold on and not go to the toilet so often. You will need to do pelvic floor muscle exercises to help you hold on.

If you wake up in the night with a strong urge to urinate, then use the toilet. As you progress you may be able to go back to sleep without getting up.

## Tips to try

- Start to avoid going 'just in case'
- Sitting down may help when you have a strong urge
- Watch the television or count back from 100 – it may take your mind off thinking about the toilet
- If able, rise up and down on tip toes or 'scrunch' toes – this makes your brain think about a physical activity rather than contracting your bladder
- Always try to avoid going to the toilet on a strong urge

## Method 1

The next time you arrive home and feel an urgent need to go to the toilet:

- Put the key in the door and before opening it, count to 60
- As you master this and the urge gets less, after counting to 60, go inside
- Go to the toilet and get ready; count to 60 before you urinate. If you leak it won't be a problem as you are over the toilet
- Continue to count to 60 each time until you do not leak any urine
- Once you have achieved this, start to extend the time from 60 seconds

## Method 2

- When you have the urge to urinate, go to the toilet and get ready
- Count to 60 before you urinate. If you leak it won't be a problem as you are over the toilet
- Count to 60 each time until you do not leak any urine
- Once you've achieved this, start to count to 60 before you go to the toilet and when you get to three, count to another 60
- Then start to extend the time from 60 seconds
- Continue extending the time until you can wait for 15 minutes without any leaking

Your anxiety about your bladder will reduce, your confidence will grow and you'll feel in control again.

## Bladder and Bowel (Continence) Service

We provide a variety of specialist continence treatments and advice for anyone with bladder or bowel problems.

You do not need to be incontinent to receive treatment and support from us. For example, you may feel that you go to the toilet too frequently, or that your bladder or bowels are limiting your lifestyle.

Your GP or any other healthcare professional can refer you to our service.

Call 0118 904 6540 10am to 2pm, Tuesday to Thursday

Email [continence@berkshire.nhs.uk](mailto:continence@berkshire.nhs.uk)



[berkshirehealthcare.nhs.uk/continence](https://berkshirehealthcare.nhs.uk/continence)